

From Low-frequency Pulsar Observations to Interstellar Holography

Pulsars make fabulous tools as probes of the interstellar medium (ISM) of our Galaxy. Their radiation is pulsed, spatially coherent and highly polarised – a combination which enables their signals to carry imprints of the ionised, turbulent and magneto-ionic properties of the media through which they propagate. At low radio frequencies (i.e. longer wavelengths), these effects are significantly magnified as a result of their strong dependencies with the observing frequency.

Multipath propagation through the ISM gives rise to a rich variety of observable effects, many of which can be meaningfully used to study the small-scale structures in the ISM. For decades, possible investigations were limited to the use of more traditional scattering and scintillation techniques, which are generally useful for a statistical characterisation of the ISM along the pulsar’s sight line. Deflected parts of the radiation may also occasionally give rise to subtle features in the secondary spectra of pulsar scintillation (e.g. parabolic arcs; Figure 1), and these can be exploited to pinpoint the location of turbulent plasma or probe any anisotropy that is present (e.g. Bhat et al. 2016, 2018). The physical origin of these arcs is an active area of research, with a multitude of recent interpretations involving hot stars or plasma sheets (Walker et al. 2017; Simard & Pen 2018; Gwinn 2019). Another notable development is the application of cyclic spectroscopy (Demorest 2011), and phase-retrieval algorithms that enable coherent de-scattering; i.e. simultaneous recovery of the pulsar’s intrinsic signal and the ISM delay structure (Walker et al. 2013).

This project will capitalise on new instrumentation and capabilities that are being developed for pulsar observations with the Murchison Widefield Array (MWA), which enable signal reconstruction at a very high time resolution (of the order of microseconds). Developing related instrumentation and the signal processing techniques, and exploiting them for novel pulsar science will form the central theme of the project. This includes accurate characterisation of signal distortion caused by the ISM (important for high-precision timing) and holographic reconstruction of the interstellar microstructure (at resolutions unattainable by other techniques). The project involves close collaboration with Manly Astrophysics.

Research Field

Observational Pulsar Astronomy

Project Suitability

Honours

Masters

PhD

Project Supervisor

Dr Ramesh Bhat

Ramesh.bhat@curtin.edu.au

Co-Supervisors

Willem van Straten (Auckland)

Mark Walker (Manley)




Fig 1: Dynamic scintillation spectrum of the millisecond pulsar J0437-4715 (left) and its secondary spectrum (right), from MWA observations (Bhat et al. 2018). Faint *parabolic arc-like* features arise from the deflected parts of pulsar’s scattered radiation. The indicated delays (\sim microseconds) will be directly measurable using the new capabilities and the advanced techniques that will be realised through this project.